



Building evidence for conservation globally

Journal of
Threatened
Taxa

Open Access

10.11609/jott.2022.14.11.22039-22206

www.threatenedtaxa.org

26 November 2022 (Online & Print)

14 (11): 22039-22206

ISSN 0974-7907 (Online)

ISSN 0974-7893 (Print)



ISSN 0974-7907 (Online); ISSN 0974-7893 (Print)

Publisher
Wildlife Information Liaison Development Society
www.wild.zooreach.org

Host
Zoo Outreach Organization
www.zooreach.org

43/2 Varadarajulu Nagar, 5th Street West, Ganapathy, Coimbatore, Tamil Nadu 641035, India
Ph: +91 9385339863 | www.threatenedtaxa.org
Email: sanjay@threatenedtaxa.org

EDITORS

Founder & Chief Editor

Dr. Sanjay Molur

Wildlife Information Liaison Development (WILD) Society & Zoo Outreach Organization (ZOO),
12 Thiruvannamalai Nagar, Saravanampatti, Coimbatore, Tamil Nadu 641035, India

Deputy Chief Editor

Dr. Neelesh Dahanukar

Noida, Uttar Pradesh, India

Managing Editor

Mr. B. Ravichandran, WILD/ZOO, Coimbatore, India

Associate Editors

Dr. Mandar Paingankar, Government Science College Gadchiroli, Maharashtra 442605, India

Dr. Ulrike Streicher, Wildlife Veterinarian, Eugene, Oregon, USA

Ms. Priyanka Iyer, ZOO/WILD, Coimbatore, Tamil Nadu 641035, India

Dr. B.A. Daniel, ZOO/WILD, Coimbatore, Tamil Nadu 641035, India

Editorial Board

Dr. Russel Mittermeier

Executive Vice Chair, Conservation International, Arlington, Virginia 22202, USA

Prof. Mewa Singh Ph.D., FASC, FNA, FNASC, FNAPsy

Ramanna Fellow and Life-Long Distinguished Professor, Biopsychology Laboratory, and Institute of Excellence, University of Mysore, Mysuru, Karnataka 570006, India; Honorary Professor, Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore; and Adjunct Professor, National Institute of Advanced Studies, Bangalore

Stephen D. Nash

Scientific Illustrator, Conservation International, Dept. of Anatomical Sciences, Health Sciences Center, T-8, Room 045, Stony Brook University, Stony Brook, NY 11794-8081, USA

Dr. Fred Pluthero

Toronto, Canada

Dr. Priya Davidar

Sigur Nature Trust, Chadapatti, Mavinhalla PO, Nilgiris, Tamil Nadu 643223, India

Dr. Martin Fisher

Senior Associate Professor, Battcock Centre for Experimental Astrophysics, Cavendish Laboratory, JJ Thomson Avenue, Cambridge CB3 0HE, UK

Dr. John Fellowes

Honorary Assistant Professor, The Kadoorie Institute, 8/F, T.T. Tsui Building, The University of Hong Kong, Pokfulam Road, Hong Kong

Prof. Dr. Mirco Solé

Universidade Estadual de Santa Cruz, Departamento de Ciências Biológicas, Vice-coordenador do Programa de Pós-Graduação em Zoologia, Rodovia Ilhéus/Itabuna, Km 16 (45662-000) Salobrinho, Ilhéus - Bahia - Brasil

Dr. Rajeev Raghavan

Professor of Taxonomy, Kerala University of Fisheries & Ocean Studies, Kochi, Kerala, India

English Editors

Mrs. Mira Bhojwani, Pune, India

Dr. Fred Pluthero, Toronto, Canada

Mr. P. Ilangoan, Chennai, India

Web Development

Mrs. Latha G. Ravikumar, ZOO/WILD, Coimbatore, India

Typesetting

Mrs. Radhika, ZOO, Coimbatore, India

Mrs. Geetha, ZOO, Coimbatore India

Fundraising/Communications

Mrs. Payal B. Molur, Coimbatore, India

Subject Editors 2019–2021

Fungi

Dr. B. Shivaraju, Bengaluru, Karnataka, India

Dr. R.K. Verma, Tropical Forest Research Institute, Jabalpur, India

Dr. Vatsavaya S. Raju, Kakatiya University, Warangal, Andhra Pradesh, India

Dr. M. Krishnappa, Jnana Sahyadri, Kuvempu University, Shimoga, Karnataka, India

Dr. K.R. Sridhar, Mangalore University, Mangalagangothri, Mangalore, Karnataka, India

Dr. Gunjan Biswas, Vidyasagar University, Midnapore, West Bengal, India

Plants

Dr. G.P. Sinha, Botanical Survey of India, Allahabad, India

Dr. N.P. Balakrishnan, Ret. Joint Director, BSI, Coimbatore, India

Dr. Shonil Bhagwat, Open University and University of Oxford, UK

Prof. D.J. Bhat, Retd. Professor, Goa University, Goa, India

Dr. Ferdinando Boero, Università del Salento, Lecce, Italy

Dr. Dale R. Calder, Royal Ontario Museum, Toronto, Ontario, Canada

Dr. Cleofas Cervancia, Univ. of Philippines Los Baños College Laguna, Philippines

Dr. F.B. Vincent Florens, University of Mauritius, Mauritius

Dr. Merlin Franco, Curtin University, Malaysia

Dr. V. Irudayaraj, St. Xavier's College, Palayamkottai, Tamil Nadu, India

Dr. B.S. Kholia, Botanical Survey of India, Gangtok, Sikkim, India

Dr. Pankaj Kumar, Kadoorie Farm and Botanic Garden Corporation, Hong Kong S.A.R., China

Dr. V. Sampath Kumar, Botanical Survey of India, Howrah, West Bengal, India

Dr. A.J. Solomon Raju, Andhra University, Visakhapatnam, India

Dr. Vijayasankar Raman, University of Mississippi, USA

Dr. B. Ravi Prasad Rao, Sri Krishnadevaraya University, Anantpur, India

Dr. K. Ravikumar, FRLHT, Bengaluru, Karnataka, India

Dr. Aparna Watve, Pune, Maharashtra, India

Dr. Qiang Liu, Xishuangbanna Tropical Botanical Garden, Yunnan, China

Dr. Noor Azhar Mohamed Shazili, Universiti Malaysia Terengganu, Kuala Terengganu, Malaysia

Dr. M.K. Vasudeva Rao, Shiv Ranjani Housing Society, Pune, Maharashtra, India

Prof. A.J. Solomon Raju, Andhra University, Visakhapatnam, India

Dr. Mandar Datar, Agharkar Research Institute, Pune, Maharashtra, India

Dr. M.K. Janarthanam, Goa University, Goa, India

Dr. K. Karthigeyan, Botanical Survey of India, India

Dr. Errol Vela, University of Montpellier, Montpellier, France

Dr. P. Lakshminarasimhan, Botanical Survey of India, Howrah, India

Dr. Larry R. Noblick, Montgomery Botanical Center, Miami, USA

Dr. K. Haridasan, Pallavur, Palakkad District, Kerala, India

Dr. Analinda Manila-Fajard, University of the Philippines Los Banos, Laguna, Philippines

Dr. P.A. Sinu, Central University of Kerala, Kasaragod, Kerala, India

Dr. Afroz Alam, Banasthali Vidyapith (accredited A grade by NAAC), Rajasthan, India

Dr. K.P. Rajesh, Zamorin's Guruvayurappan College, GA College PO, Kozhikode, Kerala, India

Dr. David E. Boufford, Harvard University Herbaria, Cambridge, MA 02138-2020, USA

Dr. Ritesh Kumar Choudhary, Agharkar Research Institute, Pune, Maharashtra, India

Dr. Navendu Page, Wildlife Institute of India, Chandrabani, Dehradun, Uttarakhand, India

Dr. Kannan C.S. Warrior, Institute of Forest Genetics and Tree Breeding, Tamil Nadu, India

Invertebrates

Dr. R.K. Avasthi, Rohtak University, Haryana, India

Dr. D.B. Bastawade, Maharashtra, India

Dr. Partha Pratim Bhattacharjee, Tripura University, Suryamaninagar, India

Dr. Kailash Chandra, Zoological Survey of India, Jabalpur, Madhya Pradesh, India

Dr. Ansie Dippenaar-Schoeman, University of Pretoria, Queenswood, South Africa

Dr. Rory Dow, National Museum of Natural History Naturalis, The Netherlands

Dr. Brian Fisher, California Academy of Sciences, USA

Dr. Richard Gallon, Ilandudno, North Wales, LL30 1UP

Dr. Hemant V. Ghate, Modern College, Pune, India

Dr. M. Monwar Hossain, Jahangirnagar University, Dhaka, Bangladesh

Mr. Jatishwor Singh Irungbam, Biology Centre CAS, Branišovská, Czech Republic.

Dr. Ian J. Kitching, Natural History Museum, Cromwell Road, UK

For Focus, Scope, Aims, and Policies, visit https://threatenedtaxa.org/index.php/JoTT/aims_scope

For Article Submission Guidelines, visit <https://threatenedtaxa.org/index.php/JoTT/about/submissions>

For Policies against Scientific Misconduct, visit https://threatenedtaxa.org/index.php/JoTT/policies_various

continued on the back inside cover

Cover: Muger Crocodile basking on the banks of Savitri River at Mahad in Maharashtra, India. © Utkarsha M. Chavan.



Paresis as a limiting factor in the reproductive efficiency of a nesting colony of *Lepidochelys olivacea* (Eschscholtz, 1829) in La Escobilla beach, Oaxaca, Mexico

Alejandra Buenrostro-Silva¹ , Jesús García-Grajales² , Petra Sánchez-Nava³
& María de Lourdes Ruíz-Gómez⁴

^{1,2} Estudiante del Programa de Doctorado en Ciencias Agropecuarias y Recursos Naturales, Universidad Autónoma del Estado de México, El Cerrillo Piedra Blancas, Toluca 50295, Estado de México.

¹ Universidad del Mar campus Puerto Escondido. Km. 2.5 carretera Federal Puerto Escondido- Sola de Vega 71980, San Pedro, Mixtepec, Oaxaca, Mexico.

^{3,4} Facultad de Ciencias, Universidad Autónoma del Estado de México. El Cerrillo Piedra Blancas, Toluca 50295, Estado de México.

¹ sba_1575@yahoo.com.mx, ² archosaurio@yahoo.com.mx (corresponding author), ³ psn@uaemex.mx, ⁴ ruiz.gomez.maria@gmail.com

Abstract: Rear flippers are crucial in the nesting process of Olive Ridley Turtles *Lepidochelys olivacea*, so any impact on them could constitute a limiting factor in reproductive efficiency. Muscle weakness of the rear legs has been observed in some nesting females on La Escobilla beach in Oaxaca state, Mexico; however, this disorder has not been sufficiently researched. The aim of this study was to identify and describe this problem in a nesting colony of *L. olivacea* in La Escobilla. We obtained the biochemical profiles of eight females with clinical signs of muscle weakness of the rear legs, that could not build the incubation chamber for their nest. In order to compare their blood characteristics, we selected eight seemingly healthy turtles that successfully built their nests, laid eggs through oviposition and covered the nest. We found no significant differences in most of the blood parameters, except for Creatinine-Kinase (CK). Female turtles with muscle weakness presented significantly higher concentrations of CK ($t = 2.1448$, d.f. = 2, $P < 0.0001$) when compared to the healthy turtles. CK is an appropriate enzyme for identifying the integrity of the muscle cell and is a muscle damage indicator. Our hypothesis is that the paresis observed in the rear legs of the female turtles in La Escobilla could be a chronic debilitation caused by a gradual exposure to biotoxins such as saxitoxins.

Keywords: Health, muscle, Olive Ridley Turtles, oviposition, rear-flippers.

Resumen: Las aletas traseras son cruciales en el proceso de anidación de la tortuga golfina *Lepidochelys olivacea*, así que cualquier impacto en ellas podría constituir un factor limitante en la eficiencia reproductiva. La debilidad muscular de las aletas traseras ha sido observada en algunas hembras anidantes de la playa La Escobilla en el estado de Oaxaca, México; sin embargo, este desorden no ha sido suficientemente investigado. El objetivo de este estudio fue identificar y describir este problema en una colonia anidante de *L. olivacea* en La Escobilla. Obtuvimos los perfiles bioquímicos de ocho hembras anidantes con signos clínicos de debilidad muscular de las aletas traseras, que impidió la construcción de la cámara de incubación para su nido. Con el fin de comparar sus características sanguíneas, seleccionamos además ocho hembras aparentemente sana que construyeron exitosamente su nido, pusieron sus huevos mediante oviposición y los cubrieron. No encontramos diferencias significativas en la mayoría de los parámetros sanguíneos, excepto por la Creatinina-Kinasa (CK). Las hembras con la debilidad muscular presentaron altas concentraciones significativas de CK ($t = 2.1448$, g.l. = 2, $P < 0.0001$) cuando fueron comparadas las tortugas aparentemente sanas. CK es una enzima apropiada para identificar la integridad de la célula muscular y es un indicador de daño muscular. Nuestra hipótesis es que la paresis observada en las patas traseras de las hembras anidantes en La Escobilla podría ser una debilitación crónica causada por una exposición gradual a biotoxinas tales como las saxitocinas.

Editor: S.R. Ganesh, Chennai Snake Park, Chennai, India.

Date of publication: 26 November 2022 (online & print)

Citation: Buenrostro-Silva, A., J. García-Grajales, P. Sánchez-Nava & M.de L. Ruíz-Gómez (2022). Paresis as a limiting factor in the reproductive efficiency of a nesting colony of *Lepidochelys olivacea* (Eschscholtz, 1829) in La Escobilla beach, Oaxaca, Mexico. *Journal of Threatened Taxa* 14(11): 22133–22138. <https://doi.org/10.11609/jott.7959.14.11.22133-22138>

Copyright: © Buenrostro-Silva et al. 2022. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use, reproduction, and distribution of this article in any medium by providing adequate credit to the author(s) and the source of publication.

Funding: None.

Competing interests: The authors declare no competing interests.

Author details, Author contributions & Acknowledgements: See end of this article.



INTRODUCTION

The Olive Ridley Turtle *Lepidochelys olivacea* (Eschscholtz, 1829) is one of the most abundant sea turtle species and classified as Vulnerable by IUCN Red List of Threatened Species (Abreu-Grobois & Plotkin 2008). In Mexico, all sea turtles are also classified as endangered species through the regulation NOM-ECOL-059-2010, and additionally, they are considered a priority species for conservation.

The Olive Ridley Turtle has a circumtropical distribution and inhabits the Atlantic, Pacific, and Indian Oceans, living mainly in the northern hemisphere (Bjorndal 1997). On the American continent, their most notable nesting beaches are located in Costa Rica (Cornelius et al. 1991; Fonseca et al. 2009), Panama (Cornelius et al. 2007; Honarvar et al. 2016), Nicaragua (Stewart 2001; Hope 2002), and Mexico (Peñaflores et al. 2000; Campbell 2007).

In Mexico, one of the world's major nesting sites for this species is La Escobilla beach in Oaxaca state; where impressively large mass synchronous nesting aggregations – called arribadas – occur (Márquez & Van Dissell 1982). Despite their importance, relatively little is known about the health status or the presence of clinical signs of a disease in the nesting colony (Mashkour et al. 2020). In addition to human activity and predation, diseases are another important factor that have contributed to the decrease in the population of sea turtles (Wallace et al. 2010).

In Olive Ridley Turtle nesting activity, rear flippers play a key role in building the nest cavity (egg chamber); hence, the depth of a nest chamber is dependent on the size of a female's rear flippers (Rusli 2019). Therefore, rear flippers are crucial in the nesting process, and any impact on them could constitute a limiting factor in reproductive efficiency.

On La Escobilla beach, various nesting turtles with muscle weakness of the rear legs have frequently been observed during the arribadas, a situation that is widely known empirically; however, the disorder has not been sufficiently researched. Until now, no studies on this subject have been performed on any nesting colony of *L. olivacea*; thus, the aim of this study was to identify and describe this problem in a nesting colony of *L. olivacea* in La Escobilla, Oaxaca, Mexico. This information will contribute to the protection of the species and allow for long-term health assessments and monitoring.

MATERIALS AND METHODS

This study was conducted on La Escobilla beach, located in the municipality of Santa Maria Tonameca on the southwestern Mexican Pacific coast (235,9041⁰N, 1451,2041⁰W). The beach is approximately 25 km long, and the Olive Ridley Turtles mainly nest along an 8 km strip at its western end.

During the development of a research project focused on assessing blood parameters in nesting Olive Ridley Turtles on La Escobilla beach, we encountered several females that displayed difficulties in making the egg chamber during the arribada events (September, October, and November) of the 2021 nesting season.

We obtained the biochemical profiles of eight females with clinical signs of muscle weakness of the rear legs that could not build the incubation chamber; as a result, after many attempts they ended up laying their eggs on the surface level of the sand (Image 1). Additionally, in order to compare their blood characteristics, we selected eight seemingly healthy turtles that successfully built their nests, laid eggs through oviposition and covered the nest. Previously, we conducted an external inspection of all the turtles evaluated to determine the existence of traumatic injuries.

Biometric data of all the turtles were taken according to the methodology described by Bolten (2000), including curved carapace length (CCL) and curved carapace width (CCW), which were measured using a flexible fiberglass tape (measurement error ± 0.5 cm, Limpus et al. 1983). The weight of each turtle was measured by suspending the turtle attached to a digital scale (MH-C 100 model, Mini Crane Scale).

Afterward, we collected blood samples from the dorsal cervical sinus, using a 1.5'-gauge needle and a 5 ml syringe and transferred the samples into vacutainers[®] containing lithium heparin as an anticoagulant (Owens & Ruiz 1980). Prior to sampling, the puncture area was cleaned. Samples were stored in refrigeration at 4 °C until laboratory processing at the Universidad del Mar (not more than eight hours after sampling).

A whole blood sample was centrifuged in an Eppendorf model 5430 centrifuge at 3,000 rpm for 10 min. We used plasma in preference to serum because in reptiles clot formation is unpredictable, changing biochemical values and occasionally producing hemolysis in the blood samples (Bolten & Bjorndal 1992). Plasma was placed posteriorly into 1.6-ml cryogenic vials. Sixteen plasma parameters were recorded using the automated blood chemistry analyzer Celercare V5 (Kabla Veterinary DX) to establish the sea turtles' health



Image 1. *Lepidochelys olivacea* with evident muscle weakness (paresis) of the rear legs laid eggs on the surface of the sand in La Escobilla beach, Oaxaca, Mexico. © Jesús García-Grajales.

profiles (Anderson et al. 2011; Espinoza-Romo et al. 2018). Additionally, we repeated the sampling process in the healthy turtles.

Analyzed parameters were divided into three groups: 1) nutrients and metabolites: Albumin (ALB; g dL⁻¹), total protein (TP), globulin (GLO), Albumin/Globulin (A/G) ratio, glucose (GLU), blood urea nitrogen (BUN), cholesterol (CHOL), creatinine (CRE), blood urea nitrogen/creatinine (BUN/CRE) ratio, total bilirubin (TBIL); 2) enzymes: amylase (AMY), Alanine aminotransferase (ALT), creatinine kinase (CK), alkaline phosphatase (ALP); and 3) electrolytes: calcium (Ca) and phosphorus (P).

Results are presented as mean, range and standard deviation (SD). Data normality and homoscedasticity were assessed using Kolmogorov-Smirnov and Levene tests, respectively. Differences in blood parameters between groups (with and with clinical signs) were assessed employing a *t*-student test. We performed all analyses using Past 4.08 statistical software (Hammer et al. 2001).

Turtle samples were collected under the SGPA/DGVS/03919/21 permit granted by SEMARNAT, Mexico.

RESULTS

Descriptive statistics of SCL, WCL, CCL, CCW, weight and blood parameters of the Olive Ridley Turtles are presented in Table 1. We found no significant differences in most of the blood parameters, except for CK.

Female turtles with muscle weakness presented significantly higher concentrations of CK ($t = 2.1448$, d.f. = 2, $p < 0.0001$) when compared with healthy turtles. During the external examination of the turtles, there was no evidence of recent traumatic injuries. However, one of the turtles with evident muscle weakness (paresis) of the rear legs laid eggs on the surface of the sand, after several failed attempts to perform the incubation chamber (Image 1).

DISCUSSION

Although Olive Ridley Turtles are the most abundant sea turtle species globally, knowledge regarding their health on mass nesting beaches remains limited. We found similar values in most blood parameters between turtles with clinical signs of paresis and seemingly healthy turtles; however, CK was highlighted as the muscle damage indicator.

Table 1. Biometric data and blood chemistry of the Olive Ridley Turtles with clinical signs of muscle weakness of the rear legs (paresis) and seemingly healthy turtles in La Escobilla beach, Oaxaca, Mexico.

	Turtles with paresis		Seemingly healthy	
	Mean (SD)	Range	Mean (SD)	Range
Biometric data				
Weight	36.2 ± 6.71	27.55 – 45.45	36.29 ± 3.56	31 – 42
CCL	65.97 ± 4.18	61.5 – 72.1	65.77 ± 2.59	63 – 69.5
CCW	69.3 ± 4.91	62.6 – 78.7	69.6 ± 2.78	65.9 – 73
Blood chemistry				
ALB	1.01 ± 0.16	0.8 – 1.3	1.15 ± 0.23	1 – 1.7
TP	3.41 ± 0.49	2.8 – 4.2	3.6 ± 0.4	3.2 – 4.5
GLO	2.4 ± 0.45	1.9 – 3.3	2.4 ± 0.2	2.2 – 2.8
A/G	0.44 ± 0.09	0.3 – 0.6	0.46 ± 0.07	0.4 – 0.6
GLU	76.6 ± 25.56	42 – 107	106.6 ± 9.4	94 – 120
BUN	17.4 ± 13.94	7.68 – 50.2	9.68 ± 1.8	7.39 – 12.6
CHOL	234.1 ± 48.06	176 – 339	244 ± 69.48	178 – 396
CRE	0.74 ± 0.23	0.46 – 0.99	0.65 ± 0.21	0.31 – 0.99
BUN/CRE	29.75 ± 33.98	11 – 109	16.5 ± 5.83	9 – 26
TBIL	0.21 ± 0.07	0.12 – 0.34	0.23 ± 0.07	0.16 – 0.39
AMY	338.5 ± 129.16	143 – 552	305.4 ± 84.9	239 – 487
ALT	2.5 ± 0.92	1 – 3	24.1 ± 9.64	11 – 45
CK *	1921.5 ± 771.79	1133 – 3143	250 ± 108.89	87 – 393
ALP	21.6 ± 5.52	13 – 31	2.875 ± 0.35	2 – 3
P	8.6 ± 1.4	6.98 – 10.55	8.2 ± 1.19	6.84 – 9.78
Ca	6.8 ± 1.74	4.9 – 10.1	5.6 ± 1.9	2 – 7.9

* Denote significant differences. CCL—Curved Carapace Length | CCW—Curved Carapace Width | ALB—Albumin | TP—Total Protein | GLO—Globulin | A/G Albumin/Globulin ratio | GLU—Glucose | BUN—Blood Urea Nitrogen | CHOL—Cholesterol | CRE—Creatinine | BUN/CRE—Blood Urea Nitrogen/Creatinine Ratio | TBIL—Total Bilirubin | AMY—Amylase | ALT—Alanine Aminotransferase | CK—Creatinine Kinase | ALP—Alkaline Phosphatase | Ca—calcium | P—Phosphorus.

CK is an appropriate enzyme for identifying the integrity of the muscle cell; and is considered a specific muscle enzyme; that is, it increases in the bloodstream when a muscle disease is present (Perrault et al. 2012; Anderson et al. 2013). With this in mind, paresis is characterized as an inability of muscles to perform their usual functions. Its physiopathology is related to the motor function of the voluntary tracks consisting of the upper and lower motor neurons, peripheral nerves, neuromuscular plate and muscle fibers, so damage to any of these structures causes a paresis that depends on the degree of the injury.

Except for Espinoza-Romo et al. (2018), there are no previous reports on CK values in *L. olivacea*. Those authors showed a CK mean of 245.3 ± 386 for *L. olivacea* in northern Sinaloa, Mexico. On the contrary, we found high CK values in turtles with obvious muscle problems of the rear legs; however, the apparently healthy females showed values close to those reported

by Espinoza-Romo et al. (2018). Because there are no papers published about muscle weakness (paresis) in *L. olivacea*, we compared our results to *Caretta caretta*, since this species has some similarities in their eating habits, sharing an analogous position in the food chain.

Previous studies on several species have reported different degrees of paresis. Jacobson et al. (2006) describes clinical signs of a neurological disorder in subadult loggerhead sea turtles (*C. caretta*) in south Florida, USA. In this study, there was no evidence of heavy metal toxicosis and organophosphate toxicosis as possible causes; instead, the clinical changes observed resulted from combined spirorchiid parasitism and possible chronic exposure to a novel toxin present in the diet of *C. caretta*.

Herrera-Galindo et al. (2015) reported several dead sea turtles (*Chelonia mydas*, *Eretmochelys imbricata*, *L. olivacea*) on the coast of Oaxaca, Mexico. This study described the presence of salps and cells of *Pyrodinium*

bahamense in the turtles' stomach contents. Later, Ley-Quióné et al. (2020) presented evidence of paralytic shellfish poisoning (PSP) causing mass mortality of sea turtles in Puerto Vallarta, Jalisco, Mexico.

PSP has been identified as the most toxic and dangerous syndrome along the Pacific coast (Sierra-Beltrán et al. 1998). Two dinoflagellate species (*Gymnodinium catenatum* and *P. bahamense*) produce saxitoxin that is associated with these PSP events (Ochoa et al. 1997; Cusik & Saylor 2013). These phenomena affect the entire trophic web, principally primary consumers, but organisms such as fish, marine mammals and sea turtles that feed on planktivorous species may also be affected (Sellner et al. 2003; Garate-Lizarraga et al. 2004).

In animals and humans, clinical signs of saxitoxin intoxication include muscular paralysis and pronounced dyspnea, which, if not promptly treated, can result in death from respiratory paralysis (Hallegraeff 1993). Although lethal doses of saxitoxins have not been defined for sea turtles or other reptiles (González-Barrientos et al. 2019), constant ingestion of this dinoflagellate species could probably provide enough toxin to produce muscle weakness of the rear legs in sea turtles.

Later, González Barrientos et al. (2019) presented evidence of abnormal clinical signs in stranded Green Sea Turtles *Chelonia mydas* that were exposed to saxitoxins and tetrodotoxins on the southern Caribbean coast of Costa Rica. Among the stranded turtles, two live green turtles exhibited extreme paresis. Saxitoxicosis in green turtles appears to have resulted from opportunistic foraging on the Caribbean Sharp-nosed Puffer *Canthigaster rostrata*.

Although specific causes of muscle weakness (paresis) in Olive Ridley Turtles remain unknown, we hypothesize that the paresis observed in the rear legs of the female turtles in La Escobilla could be a chronic debilitation due to a gradual exposure to biotoxins such as saxitoxins. It is important to mention that this debilitation could be a limiting factor in the reproductive efficiency of a nesting colony of *L. olivacea* in La Escobilla, Oaxaca; therefore, we recommended initiating a continuous monitoring program to follow the occurrence of paresis in subsequent years in order to better document its prevalence and to follow the progression of this muscle weakness among sea turtles.

REFERENCES

- Abreu-Grobois, A., & P. Plotkin (2008). *Lepidochelys olivacea*. (The IUCN Red List of Threatened Species, Database, Version 2014.2). IUCN SSC Marine Turtle Specialist Group. <https://doi.org/10.2305/IUCN.UK.2008.RLTS.T11534A3292503.en>
- Anderson, E.T., C.A. Harms, E.M. Stringer, W.M. Cluse (2011). Evaluation of hematology and serum biochemistry of cold-stunned green sea turtles (*Chelonia mydas*) in North Carolina, USA. *Journal of Zoo and Wildlife Medicine* 42(2): 247–255. <https://doi.org/10.1638/2010-0217.1>
- Anderson, E.T., V.L. Socha, J. Gardner, L. Byrd & C.A. Manire (2013). Tissue enzyme activities in the loggerhead sea turtles (*Caretta caretta*). *Journal of Zoo and Wildlife Medicine* 44(1): 62–69. <https://doi.org/10.1638/1042-7260-44.1.62>
- Bjorndal, K.A. (1997). Foraging ecology and nutrition of sea turtles, pp. 199–231. In: Lutz, P.L. & J.A. Musick (eds.). *The Biology of Sea Turtles*, CRC Press, Boca Raton, Florida, USA.
- Bolten, A.B. & K.A. Bjorndal (1992). Blood profiles for a wild population of green turtles (*Chelonia mydas*) in the Southern Bahamas: size-specific relationships. *Journal of Wildlife Diseases* 28: 407–413.
- Bolten, A.B. (2000). Técnicas para la medición de tortugas marinas, pp. 126–131. In: Eckert, K.L., K.A. Bjorndal, F.A. Abreu-Grobois, M. Donnelly (eds.). *Técnicas de investigación y manejo para la conservación de las tortugas marinas*. Grupo Especialista en Tortugas Marinas, IUCN/CSE, Washington, DC.
- Campbell, L.M. (2007). Understanding human use of Olive Ridley, implications for conservation, pp. 23–30. In: Plotkin, P.T. (ed.). *Biology and conservation of Ridley Sea turtles*. John Hopkins University Press, USA.
- Cornelius, S.E., M.A. Ulloa, J.C. Castro, M. Mata del Valle & D.C. Robinson (1991). Management of olive ridley sea turtles (*Lepidochelys olivacea*) nesting at Playas Nancite and Ostional, Costa Rica, pp. 111–135. In: Robinson, J.G. & K.H. Redfor (eds.). *Neotropical Wildlife Use and Conservation*. The University of Chicago Press, USA.
- Cornelius, S.E., R. Arauz, J. Fretey, M.H. Godfrey, R. Márquez & K. Shanker (2007). Effect of land-based harvest of *Lepidochelys*, pp. 231–251. In: Plotkin, P.T. (ed.). *Biology and conservation of Ridley Sea Turtles*, John Hopkins University Press, USA.
- Cusik K.D. & G.S. Saylor (2013). An overview on the marine neurotoxin, saxitoxin: Genetics, molecular targets, methods of detection and ecological functions. *Marine Drugs* 11: 991–1018.
- Espinoza-Romo, B.A., J.C. Sainz-Hernández, C.P. Ley-Quióné, C.E. Hart, R. Leal-Moreno, A.A. Aguirre & A.A. Zavala-Norzagaray (2018). Blood chemistry of olive ridley (*Lepidochelys olivacea*) sea turtles foraging in northern Sinaloa, Mexico. *Plos One* 13(7): e0199825. <https://doi.org/10.1371/journal.pone.0199825>
- Fonseca, L.G., G.A. Murillo, L. Guadamuz, R.M. Spinola & R.A. Valverde (2009). Downward but stable trend in the abundance of arribada olive ridley sea turtles (*Lepidochelys olivacea*) at Nancite beach, Costa Rica (1971–2007). *Chelonian Conservation and Biology* 8: 19–27.
- Garate-Lizarraga, I., J.J. Bustillos-Guzman, K. Erler, M.S. Muneton, B. Luckas & A. Tripp (2004). Paralytic shellfish toxins in the chocolate clam, *Megapitaria squalida* (Bivalvia: Veneridae), in Bahía de la Paz, Gulf of California. *Revista de Biología Tropical* 52(Suppl. 1): 133–140.
- González-Barrientos, R., G. Hernández-Mora, F. Alegre, T. Field, L. Flewelling, S. McGrath, J. Deeds, Y. Salazar, K. Rojas, E. Calvo, K. Berrocal & B. Stacy (2019). Saxitoxin poisoning in Green turtles (*Chelonia mydas*) linked to scavenging on mass mortality of Caribbean Sharpnose Puffer Fish (*Canthigaster rostrata*-Tetraodontidae). *Frontier in Veterinary Science* 6: 466. <https://doi.org/10.3389/fvets.2019.00466>
- Hallegraeff, G.M. (1993). A review of harmful algal blooms and their apparent global increase. *Phycologia* 32: 79–99.
- Hammer, Ø., D.A.T. Harper & P.D. Ryan (2001). PAST: Paleontological statistics software package for education and data analysis. *Palaentologia Electronica* 4(1): 1–9.

- Herrera-Galindo, J.E., J.F. Meraz, A. Buenrostro Silva, S. Karam, A. Mendoza & M.C. Alejo (2015). Las salpas (Thaliacea) como posible vectores de saxitoxina entre dinoflagelados y tortugas marinas. *Ciencia y Mar* 24: 41–49.
- Honarvar, S., M.C. Brodsky, E. Van Den Berghe, P. O'Connor & J. Spotila (2016). Ecology of Olive Ridley Sea Turtles at arribadas at Playa La Flor, Nicaragua. *Herpetologica* 72(4): 3003–3008.
- Hope, R.A. (2002). Wildlife harvesting, conservation and poverty: The economics of olive ridley eggs exploitation. *Environmental Conservation* 29: 375–384.
- Jacobson, E., B. Homer, B. Stacy, E. Greiner, N. Szabo, C. Chrisman, F. Origgi, S. Coberley, A. Foley, J.H. Landsberg, L. Flewelling, R.Y. Ewing, R. Moretti, S. Schaf, C. Rose, D. Mader, G.R. Harman, C.A. Manire, N.S.D. Mettee, A.P. Mizisin & G.D. Shelton (2006). Neurological disease in wild loggerhead sea turtles *Caretta caretta*. *Diseases of Aquatic Organisms* 70: 139–154.
- Ley-Quañón, C.P., C. Hart, R. Leal-Moreno, A. Martínez-López, L.A. Tello, A. Rubio, A.A. Aguirre & A.A. Zavala. (2020). Paralytic shellfish poisoning (PSP) as a cause of sea turtle mortality in Puerto Vallarta, Mexico. *Herpetological Review* 51(3): 489–494.
- Limpus, C.J., C.J. Parmenter, V. Baker & A. Fleay (1983). The crab island sea turtle rookery in north-eastern Gulf of Carpentaria. *Australian Wildlife Research* 10(1): 173–184. <https://doi.org/10.1071/WR9830173>
- Márquez, R. & H.G. Van Dissell (1985). A method for evaluating the number of massed nesting olive ridley sea turtles *Lepidochelys olivacea*, during an arribazon with comments on arribazon behavior. *Netherlands Journal of Zoology* 32: 419–425.
- Mashkour, N., K. Jones, S. Kophamel, T. Hipolito, S. Ahasan, R. Jakob-Hoff, M. Whittaker, M. Hamman, I. Bell, J. Elliman, L. Owens, C. Saladine, J.L. Crespo Picazo, B. Gardner, A.L. Loganathan, R. Bowater, E. Young, D. Robinson, W. Baverstock, D. Blyde, D. March, M. Eghbali, M. Mohammadi, D. Freggu, J. Gilliam, M. Hale, N. Nicolle, K. Spiby, D. Wrobel, M. Parga, A. Mobaraki, R. Rajakaruna, K. Hyland, M. Read & E. Ariel (2020). Disease risk analysis in sea turtles: A baseline study to inform conservation efforts. *Plos One* 15(10): e0230760. <https://doi.org/10.1371/journal.pone.0230760>
- Ochoa, J.L., A. Sánchez-Paz, A. Cruz-Villacorta & E. Núñez-Vázquez (1997). Toxic events in the northwest coastline of Mexico during 1992–1995: Origin and impact. *Hydrobiologia* 352: 195–200.
- Owens, D.W. & G.J. Ruiz (1980). New methods of obtaining blood and cerebrospinal fluid from marine turtles. *Herpetologica* 36(1): 17–20.
- Perrault, J.R., D.L. Miller, E. Eads, C. Johnson, A. Merrill, L.J. Thompson (2012). Maternal health status correlates with nest success of leatherback sea turtles (*Dermochelys coriacea*) from Florida. *Plos One* 7(2): e31841. <https://doi.org/10.1371/journal.pone.0031841>
- Peñaflores, C., J. Vasconcelos, E. Albavera, R. Márquez (2000). Twenty-five years nesting olive ridley sea turtle *Lepidochelys olivacea* in Escobilla beach, Oaxaca, Mexico, pp. 27–29. In: Abreu-Grobois, A., R. Briseño, R. Márquez & L. Sarti (eds.). Proceedings of 18th International Sea Turtle Symposium, NOAA Technical Memorandum NMFS-SEFSC-436, USA.
- Rusli, M.U. (2019). Nesting of sea turtles, pp. 1–3. In: Vonk, J. & T.K. Schakelford (eds.). *Encyclopedia of Animal Cognition and Behavior*. Springer Nature Switzerland, Germany. https://doi.org/10.1007/978-3-319-4729-6_628-1
- Sellner, K.G., G.J. Doucette & G.J. Kirkpatrick (2003). Harmful algal blooms: Causes, impacts and detection. *Journal of Indian Microbiology and Biotechnology* 30: 383–406.
- Stewart, A.Y. (2001). *Poached Modernity: Parks, People and Politics in Nicaragua, 1975–2000*. PhD dissertation, Rutgers University, USA.
- Sierra-Beltrán, A.P., E. Cruz, L.M. Del Villar, J. Cerecero & J.L. Ochoa (1998). An overview of the marine food poisoning in Mexico. *Toxicon* 36: 1493–1502.
- Wallace, B.P., A.D. DiMatteo, B.J. Hurley, E.M. Finkbeiner, A.B. Bolten & M.Y. Chaloupka (2010). Regional management units for marine turtles: a novel framework for prioritizing conservation and research across multiple scales. *Plos One* 5(12): e15465. <https://doi.org/10.1371/journal.pone.0015465>

Author details: ALEJANDRA BUENROSTRO-SILVA is currently a student of the PhD program of Ciencias Agropecuarias y Recursos Naturales of Autonomous University of the State of Mexico. Moreover, she is Professor/ Research of Universidad del Mar in the coast of Oaxaca, Mexico. Her interests include the conservation, ecology, and health of reptiles, amphibians, and mammals. JESÚS GARCÍA-GRAJALES is currently a Professor/ Research of Universidad del Mar in the coast of Oaxaca, Mexico. His interests include the tropical herpetofauna responses to anthropogenic disturbances, habitat fragmentation and conservation, ecology, and health of amphibians, reptiles and mammals. PETRA SÁNCHEZ NAVA is currently professor/scientist researcher at the Autonomous University of the State of Mexico (UAEMEX). Is responsible for the Biosustainable Systems Laboratory of the Faculty of Sciences in UAEMEX. MARIA DE LOURDES RUIZ-GÓMEZ has been a professor/scientist researcher for 10 years at the Autonomous University of the State of Mexico. Her primary interests are Animal Personalities, Behavioral Ecology, Welfare, Learning and their underlying mechanisms, especially on fish, amphibians and reptiles as models; as well as their implications on global change and conservation.

Author contributions: Development of concept and methods: ABS, JGG, PSN; fieldwork: ABS, JGG; writing: all authors; data analysis: ABS; translation: ABS-JGG.

Acknowledgements: We thank the Universidad Autónoma del Estado de México (UAEMEX) and its Division of Postgraduate Studies (PCARN) for the logistics and facilities provided. To Universidad del Mar for the facilities provided to project CUP: 21R2104. Research permits were obtained from Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT: SGPA/DGVS/03919/21). In addition, special thanks go to A.T. Rosewicz for her revisions and suggestions that improved the English of this manuscript. JG-G, PS-N, and MLRG thank the Sistema Nacional de Investigadores (SNI) for its grant.



Dr. George Mathew, Kerala Forest Research Institute, Peechi, India
Dr. John Noyes, Natural History Museum, London, UK
Dr. Albert G. Orr, Griffith University, Nathan, Australia
Dr. Sameer Padhye, Katholieke Universiteit Leuven, Belgium
Dr. Nancy van der Poorten, Toronto, Canada
Dr. Kareen Schnabel, NIWA, Wellington, New Zealand
Dr. R.M. Sharma, (Retd.) Scientist, Zoological Survey of India, Pune, India
Dr. Manju Siliwal, WILD, Coimbatore, Tamil Nadu, India
Dr. G.P. Sinha, Botanical Survey of India, Allahabad, India
Dr. K.A. Subramanian, Zoological Survey of India, New Alipore, Kolkata, India
Dr. P.M. Sureshan, Zoological Survey of India, Kozhikode, Kerala, India
Dr. R. Varatharajan, Manipur University, Imphal, Manipur, India
Dr. Eduard Vives, Museu de Ciències Naturals de Barcelona, Terrassa, Spain
Dr. James Young, Hong Kong Lepidopterists' Society, Hong Kong
Dr. R. Sundararaj, Institute of Wood Science & Technology, Bengaluru, India
Dr. M. Nithyanandan, Environmental Department, La Ala Al Kuwait Real Estate. Co. K.S.C., Kuwait
Dr. Himender Bharti, Punjabi University, Punjab, India
Mr. Purnendu Roy, London, UK
Dr. Saito Motoki, The Butterfly Society of Japan, Tokyo, Japan
Dr. Sanjay Sondhi, TITLI TRUST, Kalpavriksh, Dehradun, India
Dr. Nguyen Thi Phuong Lien, Vietnam Academy of Science and Technology, Hanoi, Vietnam
Dr. Nitin Kulkarni, Tropical Research Institute, Jabalpur, India
Dr. Robin Wen Jiang Ngiam, National Parks Board, Singapore
Dr. Lionel Monod, Natural History Museum of Geneva, Genève, Switzerland.
Dr. Asheesh Shivam, Nehru Gram Bharti University, Allahabad, India
Dr. Rosana Moreira da Rocha, Universidade Federal do Paraná, Curitiba, Brasil
Dr. Kurt R. Arnold, North Dakota State University, Saxony, Germany
Dr. James M. Carpenter, American Museum of Natural History, New York, USA
Dr. David M. Claborn, Missouri State University, Springfield, USA
Dr. Kareen Schnabel, Marine Biologist, Wellington, New Zealand
Dr. Amazonas Chagas Júnior, Universidade Federal de Mato Grosso, Cuiabá, Brasil
Mr. Moonsoon Yjoti Gogoi, Assam University, Silchar, Assam, India
Dr. Heo Chong Chin, Universiti Teknologi MARA (UiTM), Selangor, Malaysia
Dr. R.J. Shiel, University of Adelaide, SA 5005, Australia
Dr. Siddharth Kulkarni, The George Washington University, Washington, USA
Dr. Priyadarsanan Dharma Rajan, ATREE, Bengaluru, India
Dr. Phil Alderslade, CSIRO Marine And Atmospheric Research, Hobart, Australia
Dr. John E.N. Veron, Coral Reef Research, Townsville, Australia
Dr. Daniel Whitmore, State Museum of Natural History Stuttgart, Rosenstein, Germany.
Dr. Yu-Feng Hsu, National Taiwan Normal University, Taipei City, Taiwan
Dr. Keith V. Wolfe, Antioch, California, USA
Dr. Siddharth Kulkarni, The Hormiga Lab, The George Washington University, Washington, D.C., USA
Dr. Tomas Ditrich, Faculty of Education, University of South Bohemia in Ceske Budejovice, Czech Republic
Dr. Mihaly Foldvari, Natural History Museum, University of Oslo, Norway
Dr. V.P. Niyal, Wildlife Institute of India, Dehradun, Uttarakhand 248001, India
Dr. John T.D. Caleb, Zoological Survey of India, Kolkata, West Bengal, India
Dr. Priyadarsanan Dharma Rajan, Ashoka Trust for Research in Ecology and the Environment (ATREE), Royal Enclave, Bangalore, Karnataka, India

Fishes

Dr. Neelesh Dahanukar, IISER, Pune, Maharashtra, India
Dr. Topiltzin Contreras MacBeath, Universidad Autónoma del estado de Morelos, México
Dr. Heok Hee Ng, National University of Singapore, Science Drive, Singapore
Dr. Rajeev Raghavan, St. Albert's College, Kochi, Kerala, India
Dr. Robert D. Sluka, Chiltern Gateway Project, A Rocha UK, Southall, Middlesex, UK
Dr. E. Vivekanandan, Central Marine Fisheries Research Institute, Chennai, India
Dr. Davor Zanella, University of Zagreb, Zagreb, Croatia
Dr. A. Biju Kumar, University of Kerala, Thiruvananthapuram, Kerala, India
Dr. Akhilesh K.V., ICAR-Central Marine Fisheries Research Institute, Mumbai Research Centre, Mumbai, Maharashtra, India
Dr. J.A. Johnson, Wildlife Institute of India, Dehradun, Uttarakhand, India
Dr. R. Ravinesh, Gujarat Institute of Desert Ecology, Gujarat, India

Amphibians

Dr. Sushil K. Dutta, Indian Institute of Science, Bengaluru, Karnataka, India
Dr. Annemarie Ohler, Muséum national d'Histoire naturelle, Paris, France

Reptiles

Dr. Gernot Vogel, Heidelberg, Germany
Dr. Raju Vyas, Vadodara, Gujarat, India
Dr. Pritpal S. Soorae, Environment Agency, Abu Dhabi, UAE.
Prof. Dr. Wayne J. Fuller, Near East University, Mersin, Turkey
Prof. Chandrashekher U. Rivonker, Goa University, Taleigao Plateau, Goa, India
Dr. S.R. Ganesh, Chennai Snake Park, Chennai, Tamil Nadu, India
Dr. Himansu Sekhar Das, Terrestrial & Marine Biodiversity, Abu Dhabi, UAE

Birds

Dr. Hem Sagar Baral, Charles Sturt University, NSW Australia
Mr. H. Byju, Coimbatore, Tamil Nadu, India
Dr. Chris Bowden, Royal Society for the Protection of Birds, Sandy, UK
Dr. Priya Davidar, Pondicherry University, Kalapet, Puducherry, India
Dr. J.W. Duckworth, IUCN SSC, Bath, UK
Dr. Rajah Jayapal, SACON, Coimbatore, Tamil Nadu, India
Dr. Rajiv S. Kalsi, M.L.N. College, Yamuna Nagar, Haryana, India
Dr. V. Santharam, Rishi Valley Education Centre, Chittoor Dt., Andhra Pradesh, India
Dr. S. Balachandran, Bombay Natural History Society, Mumbai, India
Mr. J. Praveen, Bengaluru, India
Dr. C. Srinivasulu, Osmania University, Hyderabad, India
Dr. K.S. Gopi Sundar, International Crane Foundation, Baraboo, USA
Dr. Gombobaatar Sunde, Professor of Ornithology, Ulaanbaatar, Mongolia
Prof. Reuven Yosef, International Birding & Research Centre, Eilat, Israel
Dr. Taej Mundkur, Wetlands International, Wageningen, The Netherlands
Dr. Carol Inskipp, Bishop Auckland Co., Durham, UK
Dr. Tim Inskipp, Bishop Auckland Co., Durham, UK
Dr. V. Gokula, National College, Tiruchirappalli, Tamil Nadu, India
Dr. Arkady Lelej, Russian Academy of Sciences, Vladivostok, Russia
Dr. Simon Dowell, Science Director, Chester Zoo, UK
Dr. Mário Gabriel Santiago dos Santos, Universidade de Trás-os-Montes e Alto Douro, Quinta de Prados, Vila Real, Portugal
Dr. Grant Connette, Smithsonian Institution, Royal, VA, USA
Dr. M. Zafar-ul Islam, Prince Saud Al Faisal Wildlife Research Center, Taif, Saudi Arabia

Mammals

Dr. Giovanni Amori, CNR - Institute of Ecosystem Studies, Rome, Italy
Dr. Anwaruddin Chowdhury, Guwahati, India
Dr. David Mallon, Zoological Society of London, UK
Dr. Shomita Mukherjee, SACON, Coimbatore, Tamil Nadu, India
Dr. Angie Appel, Wild Cat Network, Germany
Dr. P.O. Nameer, Kerala Agricultural University, Thrissur, Kerala, India
Dr. Ian Redmond, UNEP Convention on Migratory Species, Lansdown, UK
Dr. Heidi S. Riddle, Riddle's Elephant and Wildlife Sanctuary, Arkansas, USA
Dr. Karin Schwartz, George Mason University, Fairfax, Virginia.
Dr. Lala A.K. Singh, Bhubaneswar, Orissa, India
Dr. Mewa Singh, Mysore University, Mysore, India
Dr. Paul Racey, University of Exeter, Devon, UK
Dr. Honnavalli N. Kumara, SACON, Anaikatty P.O., Coimbatore, Tamil Nadu, India
Dr. Nishith Dharaiya, HNG University, Patan, Gujarat, India
Dr. Spartaco Gippoliti, Socio Onorario Società Italiana per la Storia della Fauna "Giuseppe Altobello", Rome, Italy
Dr. Justus Joshua, Green Future Foundation, Tiruchirappalli, Tamil Nadu, India
Dr. H. Raghuram, The American College, Madurai, Tamil Nadu, India
Dr. Paul Bates, Harison Institute, Kent, UK
Dr. Jim Sanderson, Small Wild Cat Conservation Foundation, Hartford, USA
Dr. Dan Challenger, University of Kent, Canterbury, UK
Dr. David Mallon, Manchester Metropolitan University, Derbyshire, UK
Dr. Brian L. Cypher, California State University-Stanislaus, Bakersfield, CA
Dr. S.S. Talmale, Zoological Survey of India, Pune, Maharashtra, India
Prof. Karan Bahadur Shah, Budhanilakantha Municipality, Kathmandu, Nepal
Dr. Susan Cheyne, Borneo Nature Foundation International, Palangkaraja, Indonesia
Dr. Hemanta Kafley, Wildlife Sciences, Tarleton State University, Texas, USA

Other Disciplines

Dr. Aniruddha Belsare, Columbia MO 65203, USA (Veterinary)
Dr. Mandar S. Paingankar, University of Pune, Pune, Maharashtra, India (Molecular)
Dr. Jack Tordoff, Critical Ecosystem Partnership Fund, Arlington, USA (Communities)
Dr. Ulrike Streicher, University of Oregon, Eugene, USA (Veterinary)
Dr. Hari Balasubramanian, EcoAdvisors, Nova Scotia, Canada (Communities)
Dr. Rayanna Hellem Santos Bezerra, Universidade Federal de Sergipe, São Cristóvão, Brazil
Dr. Jamie R. Wood, Landcare Research, Canterbury, New Zealand
Dr. Wendy Collinson-Jonker, Endangered Wildlife Trust, Gauteng, South Africa
Dr. Rajeshkumar G. Jani, Anand Agricultural University, Anand, Gujarat, India
Dr. O.N. Tiwari, Senior Scientist, ICAR-Indian Agricultural Research Institute (IARI), New Delhi, India
Dr. L.D. Singla, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana, India
Dr. Rupika S. Rajakaruna, University of Peradeniya, Peradeniya, Sri Lanka
Dr. Bahar Baviskar, Wild-CER, Nagpur, Maharashtra 440013, India

Reviewers 2019–2021

Due to paucity of space, the list of reviewers for 2018–2020 is available online.

The opinions expressed by the authors do not reflect the views of the Journal of Threatened Taxa, Wildlife Information Liaison Development Society, Zoo Outreach Organization, or any of the partners. The journal, the publisher, the host, and the partners are not responsible for the accuracy of the political boundaries shown in the maps by the authors.

Journal of Threatened Taxa is indexed/abstracted in Bibliography of Systematic Mycology, Biological Abstracts, BIOSIS Previews, CAB Abstracts, EBSCO, Google Scholar, Index Copernicus, Index Fungorum, JournalSeek, National Academy of Agricultural Sciences, NewJour, OCLC WorldCat, SCOPUS, Stanford University Libraries, Virtual Library of Biology, Zoological Records.

NAAS rating (India) 5.64

Print copies of the Journal are available at cost. Write to:
The Managing Editor, JoTT,
c/o Wildlife Information Liaison Development Society,
43/2 Varadarajulu Nagar, 5th Street West, Ganapathy, Coimbatore,
Tamil Nadu 641035, India
ravi@threatenedtaxa.org



www.threatenedtaxa.org

OPEN ACCESS



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

November 2022 | Vol. 14 | No. 11 | Pages: 22039-22206

Date of Publication: 26 November 2022 (Online & Print)

DOI: 10.11609/jott.2022.14.11.22039-22206

Communications

New records of pteridophytes in Mount Matutum Protected Landscape, South Central Mindanao, Philippines with notes on its economic value and conservation status

– Christine Dawn Galope-Obemio, Inocencio E. Buot Jr. & Maria Celeste Banaticla-Hilario, Pp. 22039–22057

Some threatened woody plant species recorded from forests over limestone of the Philippines

– Inocencio E. Buot Jr., Marne G. Origenes, Ren Divien R. Obeña, Elaine Loreen C. Villanueva & Marjorie D. delos Angeles, Pp. 22058–22079

Status of mangrove forest in Timaco Mangrove Swamp, Cotabato City, Philippines

– Cherie Cano-Mangaoang, Zandra Caderon Amino & Baingan Brahim Mastur, Pp. 22080–22085

A comparative analysis of the past and present occurrences of some species of *Paphiopedilum* (Orchidaceae) in northeastern India using MaxEnt and GeoCAT

– Debonina Dutta & Aparajita De, Pp. 22086–22097

Foraging activity and breeding system of *Avicennia officinalis* L. (Avicenniaceae) in Kerala, India

– K. Vinaya & C.F. Binoy, Pp. 22098–22104

Diversity patterns and seasonality of hawkmoths (Lepidoptera: Sphingidae) from northern Western Ghats of Maharashtra, India

– Aditi Sunil Shere-Kharwar, Sujata M. Magdum, G.D. Khedkar & Supriya Singh Gupta, Pp. 22105–22117

Population trends of Mugger Crocodile and human-crocodile interactions along the Savitri River at Mahad, Maharashtra, India

– Utkarsha Manish Chavan & Manoj Ramakant Borkar, Pp. 22118–22132

Paresis as a limiting factor in the reproductive efficiency of a nesting colony of *Lepidochelys olivacea* (Eschscholtz, 1829) in La Escobilla beach, Oaxaca, Mexico

– Alejandra Buenrostro-Silva, Jesús García-Grajales, Petra Sánchez-Nava & María de Lourdes Ruíz-Gómez, Pp. 22133–22138

Notes on the nesting and foraging behaviours of the Common Coot *Fulica atra* in the wetlands of Viluppuram District, Tamil Nadu, India

– M. Pandian, Pp. 22139–22147

Population abundance and threats to Black-headed Ibis *Threskiornis melanocephalus* and Red-naped Ibis *Pseudibis papillosa* at study sites in Jhajjar district, Haryana, India

– Anjali & Sarita Rana, Pp. 22148–22155

Crop raiding and livestock predation by wildlife in Khaptad National Park, Nepal

– Ashish Bashyal, Shyam Sharma, Narayan Koirala, Nischal Shrestha, Nischit Aryal, Bhupendra Prasad Yadav & Sandeep Shrestha, Pp. 22156–22163

Review

An annotated checklist of odonates of Amboli-Chaukul-Parpoli region showing new records for the Maharashtra State, India with updated state checklist

– Dattaprasad Sawant, Hemant Ogale & Rakesh Mahadev Deulkar, Pp. 22164–22178

Short Communications

The new addition of Blue Pimpernel of Primulaceae to the state flora of Assam, India

– Sushmita Kalita, Barnali Das & Namita Nath, Pp. 22179–22183

A new species of genus *Neocerura* Matsumura, 1929 (Notodontidae: Lepidoptera) from India

– Amritpal Singh Kaleka & Rishi Kumar, Pp. 22184–22189

Rediscovery of an interesting preying mantis *Deiphobella laticeps* (Mantodea: Rivetiniidae) from Maharashtra, India

– Gauri Sathaye, Sachin Ranade & Hemant V. Ghate, Pp. 22190–22194

Camera trapping records confirm the presence of the elusive Spotted Linsang *Prionodon pardicolor* (Mammalia: Carnivora: Prionodontidae) in Murlen National Park (Mizoram, India)

– Amit Kumar Bal & Anthony J. Giordano, Pp. 22195–22200

Notes

First sighting record of the Orange-breasted Green-Pigeon *Treron bicinctus* (Aves: Columbiformes: Columbidae) from Chittaranjan, West Bengal, India

– Shahbaz Ahmed Khan, Nazneen Zehra & Jamal Ahmad Khan, Pp. 22201–22202

Book Reviews

Decoding a group of winged migrants!

– Review by Priyanka Iyer, Pp. 22203–22204

First steps of citizen science programs in India

– Review by Aishwarya S. Kumar & Lakshmi Nair, Pp. 22205–22206

Publisher & Host

